

CUET – 2023

Mathematics

Mock Paper – 1 (Solution)

Time: 45 min

Maximum Marks: 200

General Instructions:

- (i) Total duration of **Mathematics** Paper is **45 min**.
- (ii) You have to attempt 40 questions out of 50 in each Domain subjects.
- (iii) All the questions provided are in MCQ format and have only single correct option.
- (iv) Each question carries 5 marks. For each correct response, the candidate will get 5 marks. For each incorrect response, 1 mark will be deducted from the total score.


1. If $A = \{1, 2, 3, 4\}$ and $B = \{5, 6, 7, 8\}$, then the function which is one-one and onto is

- (a) $f_1 = \{(1, 5), (2, 7), (3, 8), (4, 6)\}$
- (b) $f_2 = \{(1, 6), (2, 8), (3, 8), (4, 5)\}$
- (c) $f_3 = \{(1, 5), (2, 7), (3, 8), (4, 5)\}$
- (d) $f_4 = \{(1, 8), (2, 7), (3, 6), (4, 7)\}$

Correct answer – a: $f_1 = \{(1, 5), (2, 7), (3, 8), (4, 6)\}$

Explanation:

The function $f_1: A \rightarrow B$ is given below:



In this function, the second entry in each ordered pair is unique and the set of second entries of the ordered pairs is the set B.

So, the function f_1 is one-one and onto.

2. The cosine function can be restricted to any interval of the type____, for its inverse to exist.

- (a) $[n\pi, (n+1)\pi]$
- (b) $[n\pi/2, (n+1)\pi/2]$
- (c) $(n\pi, (n+1)\pi)$
- (d) $(n\pi/2, (n+1)\pi/2)$

Correct answer – c: $(n\pi, (n+1)\pi)$

Explanation:

The cosine function restricted to any of the intervals $[-\pi, 0]$, $[0, \pi]$, $[\pi, 2\pi]$ etc., is bijective with range $[-1, 1]$.

We can, therefore, define the inverse of cosine function in each of these intervals.

The general form of these intervals is $(n\pi, (n+1)\pi)$.

3. A square matrix A is called an orthogonal matrix if

- (a) $AA^T = A^TA = I$
- (b) $AI = A$
- (c) $A^2 = I$
- (d) $A(A^T)^T = I$

Correct answer – a: $AA^T = A^TA = I$

Explanation:

A square matrix A is called an orthogonal matrix when $AA^T = A^TA = I$

4. If $A = \begin{bmatrix} \cos x & \sin x \\ -\sin x & \cos x \end{bmatrix}$ and $A(\text{adj } A) = k \begin{bmatrix} 5 & 0 \\ 0 & 5 \end{bmatrix}$, then the value of k is:

- (a) $\frac{1}{3}$
- (b) $\frac{1}{4}$
- (c) $\frac{1}{5}$
- (d) $\frac{1}{6}$

Correct answer – c: $\frac{1}{5}$

Explanation:

We know that, $A(\text{adj } A) = |A|I$

If $A = \begin{bmatrix} \cos x & \sin x \\ -\sin x & \cos x \end{bmatrix}$, then

$$|A| = \cos^2 x + \sin^2 x = 1$$

$$\therefore A (\text{adj } A) = I$$

It is given that,

$$A (\text{adj } A) = k \begin{bmatrix} 5 & 0 \\ 0 & 5 \end{bmatrix} = 5k \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$$

$$\Rightarrow A (\text{adj } A) = 5k I$$

$$\Rightarrow 5k = 1 \Rightarrow k = \frac{1}{5}$$

5. If $A = \begin{bmatrix} 2 & -5 \\ -3 & 1 \end{bmatrix}$, then $\text{adj } (4A^2 + 9A)$ is equal to:

(a) $\begin{bmatrix} 94 & 105 \\ 63 & 73 \end{bmatrix}$

(b) $\begin{bmatrix} 73 & 63 \\ 105 & 94 \end{bmatrix}$

(c) $\begin{bmatrix} 105 & 73 \\ 63 & 94 \end{bmatrix}$

(d) $\begin{bmatrix} 73 & 105 \\ 63 & 94 \end{bmatrix}$

Correct answer - d: $\begin{bmatrix} 73 & 105 \\ 63 & 94 \end{bmatrix}$

Explanation:

$$\text{Given: } A = \begin{bmatrix} 2 & -5 \\ -3 & 1 \end{bmatrix}$$

$$A^2 = \begin{bmatrix} 2 & -5 \\ -3 & 1 \end{bmatrix} \begin{bmatrix} 2 & -5 \\ -3 & 1 \end{bmatrix} = \begin{bmatrix} 4 + 15 & -10 - 5 \\ -6 - 3 & 15 + 1 \end{bmatrix} = \begin{bmatrix} 19 & -15 \\ -9 & 16 \end{bmatrix}$$

$$\therefore 4A^2 = 4 \begin{bmatrix} 19 & -15 \\ -9 & 16 \end{bmatrix} = \begin{bmatrix} 76 & -60 \\ -36 & 64 \end{bmatrix}$$

$$9A = \begin{bmatrix} 18 & -45 \\ -27 & 9 \end{bmatrix}$$

$$4A^2 + 9A = \begin{bmatrix} 76 & -60 \\ -36 & 64 \end{bmatrix} + \begin{bmatrix} 18 & -45 \\ -27 & 9 \end{bmatrix} = \begin{bmatrix} 94 & -105 \\ -63 & 73 \end{bmatrix}$$

$$\text{adj } (4A^2 + 9A) = \begin{bmatrix} 73 & 63 \\ 105 & 94 \end{bmatrix}^T = \begin{bmatrix} 73 & 105 \\ 63 & 94 \end{bmatrix}$$

6. If $f: \mathbb{R} \rightarrow \mathbb{R}$ and $f(x) = x$, then $f \circ f = ?$

(a) \mathbb{R}

(b) x

(c) $2x$

(d) $3x$

Correct answer – b: x

Explanation:

$$(f \circ f)(x) = f\{f(x)\} = f\{x\} = x$$

7. If $f(x) = |x - 2|$, then at $x = 2$, $f'(x)$ is

- (a) Continuous but not differentiable
- (b) Differentiable but not continuous
- (c) Continuous and differentiable both
- (d) Neither continuous nor differentiable

Correct answer – a: Continuous but not differentiable

Explanation:

$$f'(x) = -1 \text{ at } x < 2$$

$$f'(x) = 1 \text{ at } x > 2$$

Therefore, not differentiable but continuous as it is a composition of two functions i.e., polynomial and modulus.

8. Find the second derivative of $e^x \cos x$.

- (a) $-e^x \sin x$
- (b) $-2e^x \cos x$
- (c) $-2e^x \sin x$
- (d) $e^x(\sin x + \cos x)$

Correct answer – c: $-2e^x \sin x$

Explanation:

$$\text{Let } y = e^x \cos x$$

$$\frac{dy}{dx} = e^x(\cos x - \sin x)$$

$$\frac{d^2y}{dx^2} = e^x(\cos x - \sin x) + e^x(-\sin x - \cos x)$$

$$= -2e^x \sin x$$

9. What is the maximum number of different elements needed to write a skew symmetric matrix of order n ?

- (a) n^2
- (b) n
- (c) $n^2 - n$
- (d) $n^2 - n + 1$

Correct answer – d: $n^2 - n + 1$

Explanation:

To write a square matrix of order n , we need n^2 elements.

In a skew symmetric matrix, all the diagonal elements are zeros.

So, we need only $n^2 - n + 1$ elements to write a skew symmetric matrix of order n .

10. Evaluate: $\cos(\tan^{-1} x)$

- (a) $\frac{1}{1-x^2}$
- (b) $\frac{1}{1+x^2}$
- (c) $\frac{1}{\sqrt{1-x^2}}$
- (d) $\frac{1}{\sqrt{1+x^2}}$

Correct answer – d: $\frac{1}{\sqrt{1+x^2}}$

Explanation:

We have,

$$\cos(\tan^{-1} x) = \cos\left(\cos^{-1} \frac{1}{\sqrt{1+x^2}}\right) = \frac{1}{\sqrt{1+x^2}}$$

11. To check whether matrix B is an inverse of matrix A , we need to check

- (a) $AB^{-1} = I$
- (b) $BA^{-1} = I$
- (c) $AB = BA = I$
- (d) Either $AB = I$ or $BA = I$

Correct answer – c: $AB = BA = I$

Explanation:

If A is a square matrix of order m , and if there exists another square matrix B of the same order m , such that $AB = BA = I$, then B is called the inverse matrix of A and it is denoted by A^{-1} .

In that case A is said to be invertible.

12. The greatest integer function is:

- (a) continuous everywhere
- (b) discontinuous everywhere
- (c) continuous except at the integral values of x
- (d) discontinuous except at end points

Correct answer – c: continuous except at the integral values of x

Explanation:

The graph of greatest integer function $[x]$ breaks at integral values of x .

Thus, it is continuous everywhere except at the integral values of x .

13. A real function f is said to be continuous if it is continuous at every point in

- (a) any interval of real numbers
- (b) $[-\infty, \infty]$
- (c) the range of f
- (d) the domain of f

Correct answer – d: the domain of f

Explanation:

A real function f is said to be continuous if it is continuous at every point in the domain of f .

14. The area enclosed between the lines $x = 2$ and $x = 7$ is

- (a) 7 units
- (b) 5 units
- (c) 2 units
- (d) ∞

Correct answer – d: ∞

Explanation:


Area between two parallel lines is infinite.

15. Area of the region bounded by the curve $y^2 = 2y - x$ and y-axis is:

- (a) 3 sq. units
- (b) 4 sq. units
- (c) $3/4$ sq. units
- (d) $4/3$ sq. units

Correct answer – d: $4/3$ sq. units

Explanation:



Putting $x = 0$ in $y^2 = 2y - x$, $y = 0$ and $y = 2$

$$Area = \int_0^2 x dy$$

$$\Rightarrow A = \int_0^2 (2y - y^2) dy$$

$$\Rightarrow A = \left[y^2 - \frac{y^3}{3} \right]_0^2 = 4 - \frac{8}{3} = \frac{4}{3}$$

$$Area = \frac{4}{3} \text{ sq. units}$$

16. The first order linear differential equation, where y is independent and x is dependent variable, is given by:

(a) $\frac{dy}{dx} + P(x)y = Q(x)$

(b) $\frac{dx}{dy} + P(x)y = Q(x)$

(c) $\frac{dx}{dy} + P(y)x = Q(x)$

(d) $\frac{dy}{dx} + P(y)x = Q(y)$

Correct answer - d: $\frac{dy}{dx} + P(y)x = Q(y)$

Explanation:

The first order linear differential equation, where y is independent and x is dependent variable, is:

$$\frac{dx}{dy} + P(y).x = Q(y)$$

Here, P(y) and Q(y) are the functions of y.

17. Which of the following is not a linear differential equation?

(a) $\frac{dy}{dx} + (1 + x^2)y = (1 + x^2)^2$

(b) $\frac{dy}{dx} + (1 + xy)y = x^2 + 2$

(c) $\frac{dy}{dx} + 1 + 2y = 4x$

(d) $\frac{dy}{dx} + \left(\frac{\cos x}{\sin x}\right) = \tan x$

Correct answer - b: $\frac{dy}{dx} + (1 + xy)y = x^2 + 2$

Explanation:

$\frac{dy}{dx} + (1 + xy)y = x^2 + 2$ is not a linear differential equation.

18. If O be the origin and $P_1(x_1, y_1, z_1)$ & $P_2(x_2, y_2, z_2)$ are two points, then the vector joining the points P_1 and P_2 is the vector P_1P_2 given by

(a) $OP_1 + OP_2$

(b) $OP_2 - OP_1$

(c) $OP_1.OP_2$

(d) $OP_1 - OP_2$

Correct answer - b: $OP_2 - OP_1$

Explanation:

$$OP_1 + P_1P_2 = OP_2 \text{ or } P_1P_2 = OP_2 - OP_1$$

19. What is the additive identity of a vector?

- (a) zero vector
- (b) unit vector
- (c) The vector itself
- (d) Negative of the vector

Correct answer – a: zero vector

Explanation:

$$\vec{a} + \vec{0} = \vec{0} + \vec{a} = \vec{a}$$

Therefore, zero vector is the additive identity for a vector.

20. Area bounded by the curve $y = \cos x$ between $x = 0$ and $x = 2\pi$ is

- (a) 0
- (b) 1 square unit
- (c) 2 square units
- (d) 4 square units

Correct answer – d: 4 square units

Explanation:

The required area is given by

$$\begin{aligned} A &= \int_0^{2\pi} \cos x \, dx \\ &= \int_0^{\frac{\pi}{2}} \cos x \, dx + \int_{\frac{\pi}{2}}^{\frac{3\pi}{2}} \cos x \, dx + \int_{\frac{3\pi}{2}}^{2\pi} \cos x \, dx \\ &= [\sin x]_0^{\frac{\pi}{2}} + [\sin x]_{\frac{\pi}{2}}^{\frac{3\pi}{2}} + [\sin x]_{\frac{3\pi}{2}}^{2\pi} \\ &= 1 + 2 + 1 \dots \text{(Since area can't be -ve)} \\ &= 4 \text{ sq. units} \end{aligned}$$

21. The distance of the point $(2, 3, -5)$ from the plane $x + 2y - 2z = 9$ is:

- (a) 2 units
- (b) $3/2$ units
- (c) 3 units
- (d) $10/3$ units

Correct answer – c: 3 units

Explanation:

The distance of the point $(2, 3, -5)$ from the plane $x + 2y - 2z = 9$ is:

$$d = \frac{2(1) + 2(3) - 2(-5) - 9}{\sqrt{1+4+4}} = \frac{2+6+10-9}{\sqrt{9}} = \frac{9}{3} = 3$$

22. The Cartesian equation of the line passing through the points (a, b, c) and (a', b', c') is:

(a) $\frac{x-a}{a'-a} = \frac{y-b}{b'-b} = \frac{z-c}{c'-c}$

(b) $\frac{x-a}{a'+a} = \frac{y-b}{b'+b} = \frac{z-c}{c'+c}$

(c) $\frac{x+a}{a'-a} = \frac{y+b}{b'-b} = \frac{z+c}{c'-c}$

(d) $\frac{x+a}{a'+a} = \frac{y+b}{b'+b} = \frac{z+c}{c'+c}$

Correct answer – a: $\frac{x-a}{a'-a} = \frac{y-b}{b'-b} = \frac{z-c}{c'-c}$

Explanation:

The Cartesian equation of the line passing through the points (a, b, c) and (a', b', c') is:

$$\frac{x-a}{a'-a} = \frac{y-b}{b'-b} = \frac{z-c}{c'-c}$$

23. The length of the perpendicular from the origin to the plane $3x + 2y - 6z = 21$ is:

(a) 7

(b) 14

(c) 3

(d) 21

Correct answer – c: 3

Explanation:

The perpendicular distance from the origin to the plane is

$$= \frac{|3(0) + 2(0) - 6(0) - 21|}{\sqrt{(3)^2 + (2)^2 + (-6)^2}} = \frac{|-21|}{\sqrt{49}} = \frac{21}{7} = 3$$

24. Objective function of an LPP is

- (a) a constraint
- (b) a function to be optimized
- (c) a relation between the variables
- (d) equation in a line

Correct answer – b: a function to be optimized

Explanation:

Objective function of an LPP is a function to be optimized.


25. The solution set of the inequation $2x + y > 5$ is

- (a) half plane that contains the origin
- (b) open half plane not containing the origin
- (c) whole xy -plane except the points lying on the line $2x + y = 5$
- (d) points on line $2x + y = 5$

Correct answer – b: open half plane not containing the origin

Explanation:

Consider the following graph.



Thus, the graph of the given equation is the open half plane not containing the origin.

26. Let $Z = ax + by$ be a linear objective function, then variables x and y are called _____ variables.

- (a) dependent
- (b) independent
- (c) decision
- (d) continuous

Correct answer – c: Decision

Explanation:

If $Z = ax + by$ is a linear objective function, then variables x and y are called decision variables.

27. Vectors that may subject to its parallel displacement without changing its magnitude and direction are called _____.

- (a) free vectors
- (b) co-initial vectors
- (c) parallel vectors
- (d) collinear vectors

Correct answer – a: Free vectors

Explanation:

Vectors that may subject to its parallel displacement without changing its magnitude and direction are called free vectors.

28. Two or more vectors having the same initial point are called

- (a) unit vectors
- (b) zero vectors
- (c) co-initial vectors
- (d) co-terminus vectors

Correct answer – c: co-initial vectors

Explanation:

Two or more vectors having the same initial point are called co-initial vectors.

29. The order and degree of the differential equation: $(y'')^2 + (y'')^3 + (y')^4 + y^5 = 0$ are respectively:

- (a) 2 and 4
- (b) 2 and 3
- (c) 2 and 5
- (d) 3 and 5

Correct answer – b: 2 and 3

Explanation:

Since, the highest differential coefficient of the equation $(y'')^2 + (y'')^3 + (y')^4 + y^5 = 0$ is y'' and power of y'' is 3.

Therefore, order of the equation is 2 and degree is 3.

30. The solution of the below differential equation is:

$$\frac{dy}{dx} = x \log x$$

(a) $y = \frac{x^2}{2} \log x - \frac{x^2}{4} + c$

(b) $y = \frac{x^2}{2} \log x + \frac{x^2}{4} + c$

(c) $2y = x^2(\log x + 1) + c$

(d) $y = x^2(\log x + 1) + c$

Correct answer - a: $y = \frac{x^2}{2} \log x - \frac{x^2}{4} + c$

Explanation:

$$\frac{dy}{dx} = x \log x \Rightarrow dy = x \log x \, dx$$

$$\Rightarrow \int dy = \int x \log x \, dx$$

$$\Rightarrow y = \log x \cdot \frac{x^2}{2} - \int \frac{1}{x} \cdot \frac{x^2}{2} \, dx + c$$

$$\Rightarrow y = \frac{x^2}{2} \log x - \frac{x^2}{4} + c$$

31. The curves $x^2 + y^2 = 16$ and $y^2 = 6x$ intersects at

(a) $(2, 2\sqrt{3})$

(b) $(0, 2\sqrt{3})$

(c) $(2, 0)$

(d) $(0, 2)$

Correct answer - a: $(2, 2\sqrt{3})$

Explanation:

$$x^2 + y^2 = 16; y^2 = 6x$$

The points of intersection of the two curves $x^2 + y^2 = 16$ and $y^2 = 6x$

$$x^2 + 6x = 16 \Rightarrow x^2 + 6x - 16 = 0 \Rightarrow (x+8)(x-2) = 0 \Rightarrow x = -8, 2$$

$$\text{But } x \text{ is non negative} \Rightarrow x = 2 \Rightarrow y^2 = 6(2) = 12$$

$$\Rightarrow y = \sqrt{12} = 2\sqrt{3}$$

\therefore The points of intersection of the two curves is $(2, 2\sqrt{3})$

32. If $\int_0^{\alpha} \frac{1}{1+4x^2} dx = \frac{\pi}{8}$, the value of α is

- (a) $\frac{1}{2} \tan \frac{\pi}{8}$
- (b) $\frac{1}{2}$
- (c) 1
- (d) $\frac{1}{2\sqrt{2}}$

Correct answer – b: $\frac{1}{2}$

Explanation:

$$\text{Given: } \int_0^{\alpha} \frac{1}{1+4x^2} dx = \frac{\pi}{8}$$

$$\Rightarrow \frac{1}{2} [\tan^{-1}(2x)]_0^{\alpha} = \frac{\pi}{8}$$

$$\Rightarrow \tan^{-1}(2\alpha) = \frac{\pi}{4}$$

$$\Rightarrow 2\alpha = \tan \frac{\pi}{4} \Rightarrow \alpha = \frac{1}{2}$$

33. The integration of the function $\log x$ is:

- (a) $x(\log x - 1) + C$
- (b) $(x \log x - 1) + C$
- (c) $x(\log x + 1) + C$
- (d) $(x \log x + 1) + C$

Correct answer – a: $x(\log x - 1) + C$

Explanation:

$$\begin{aligned} \int \log x \, dx &= \int \log x (1) \, dx = \log x \cdot x - \int \frac{1}{x} \cdot x \, dx \\ &= x \log x - x + C = x(\log x - 1) + C \end{aligned}$$

34. The value of definite integral depends on

- (a) the function and the interval
- (b) the interval and the variable of integration
- (c) the function and the variable of integration
- (d) the function, the interval and the variable of integration

Correct answer – a: the function and the interval

Explanation:

Since in definite integrals, the variable of integration is a dummy variable. The value of definite integral depends only on the function and the interval not on the variable of integration.

35. Integration of $\sec x$ is:

- (a) $\log |\sec x| + c$
- (b) $\log |\tan x| + c$
- (c) $\log |\sec x + \tan x| + c$
- (d) $\frac{1}{\log(\sec x)} + c$

Correct answer – c: $\log |\sec x + \tan x| + c$

Explanation:

$$\text{Let } \int \sec x dx = I = \int \frac{\sec x (\sec x + \tan x)}{(\sec x + \tan x)} dx$$

$$\text{Put } (\sec x + \tan x) = t \Rightarrow (\sec x \tan x + \sec^2 x) dx = dt$$

$$\Rightarrow \sec x (\sec x + \tan x) dx = dt$$

$$\therefore I = \int \frac{1}{t} dt = \log |t| + c = \log |\sec x + \tan x| + c$$

36. Geometrically Rolle's theorem ensures that there is at least one point on the curve $f(x)$, whose abscissa lies in (a, b) at which the tangent is


- (a) parallel to the y-axis
- (b) parallel to the x-axis
- (c) parallel to the line $y = x$
- (d) parallel to the line joining the end points of the curve

Correct answer – b: parallel to the x-axis

Explanation:

Rolle's Theorem states that Let $f : [a, b] \rightarrow \mathbb{R}$ be continuous on $[a, b]$ and be differentiable on (a, b) , such that $f(a) = f(b)$ where a and b are some real numbers. Then there exists some c in (a, b) such that $f'(c) = 0$.

The geometrical meaning of the statement is that when $f(x)$ satisfies all relevant conditions, then, there is atleast one point on the curve $f(x)$, whose abscissa lies in (a, b) at which the tangent is parallel to the x axis



37. If order of the matrix A is 2×3 , of matrix B is 3×2 , and of matrix C is 3×3 , then which one of the following is not defined?

- (a) $C(A + B')$
- (b) $C(A + B')'$
- (c) BAC
- (d) $CB + A'$

Correct answer – a: $C(A + B')$

Explanation:

$A + B'$ is a matrix of order 2×3

Since, C is a matrix of order 3×3 .

So, $C(A + B')$ is not defined.

38. If A is a matrix of order $m \times n$ and B is a matrix of order $l \times p$. The product AB of two matrices is defined if,

- (a) $n = p$
- (b) $m = p$
- (c) $n = l$
- (d) $m = l$

Correct answer – c: $n = l$

Explanation:

The product of two matrices A and B is defined if the number of columns of A is equal to the number of rows of B.

So here, AB is defined if $n = l$.

39. If $x + y + z = xyz$, then $\tan^{-1}x + \tan^{-1}y + \tan^{-1}z =$

- (a) π
- (b) $\pi/2$
- (c) 1
- (d) $\tan^{-1}(xyz)$

Correct answer – a: π

Explanation:

$$\tan^{-1}x + \tan^{-1}y + \tan^{-1}z = \tan^{-1}\left(\frac{x + y + z - xyz}{1 - xy - yz - zx}\right) = \tan^{-1}0 = \pi$$

40. If $A = N \times N$ and $*$ be any binary operation on A defined by $(a, b) * (c, d) = (a + c, b + d)$, then the binary operation is

- (a) commutative
- (b) associative
- (c) commutative and associative
- (d) commutative but not associative

Correct answer – c: commutative and associative

Explanation:

$(a, b) * (c, d) = (a + c, b + d)$ and $(c, d) * (a, b) = (c + a, d + b) = (a + c, b + d) = (a, b) * (c, d)$ So, $*$ is commutative.

$$(a, b) * [(c, d) * (e, f)] = (a, b) * (c + e, d + f) = (a + c + e, b + d + f)$$

$$\text{Also, } [(a, b) * (c, d)] * (e, f) = (a + c, b + d) * (e, f) = (a + c + e, b + d + f).$$

So, $*$ is associative also.

Hence, $*$ is commutative and associative.

41. What is the probability of $(A \cup R) \cap S$?

- (a) $\frac{1}{6}$
- (b) $\frac{2}{6}$
- (c) $\frac{3}{6}$
- (d) $\frac{1}{3}$

Correct Option - c: $\frac{3}{6}$

Explanation:

$(A \cup R) \cap S = \{1, 2, 5\}$ and Sample Space = $\{1, 2, 3, 4, 5, 6\}$

$$P((A \cup R) \cap S) = \frac{n((A \cup R) \cap S)}{n(\text{Sample space})} = \frac{3}{6}$$

42. The value of $P(A | R)$ is

- (a) $\frac{1}{6}$
- (b) $\frac{2}{6}$
- (c) $\frac{3}{6}$
- (d) $\frac{1}{3}$

Correct Option - d: $\frac{1}{3}$

Explanation:

Here, sample space = $\{1, 2, 3, 4, 5, 6\}$

$A \cap R = \{5\}$, $R \cap S = \{2, 5\}$, $A \cap S = \{1, 5\}$, $A \cap R \cap S = \{5\}$, $(A \cup R) \cap S = \{1, 2, 5\}$

$$P(A) = \frac{2}{6} = \frac{1}{3}, P(R) = \frac{3}{6} = \frac{1}{2}, P(S) = \frac{3}{6} = \frac{1}{2}$$

$$\text{Also, } P(A \cap R) = \frac{1}{6}, P(R \cap S) = \frac{2}{6}, P(A \cap S) = \frac{2}{6}$$

$$P(A \cap R \cap S) = \frac{1}{6} \text{ and } P((A \cup R) \cap S) = \frac{3}{6}$$

$$P(A|R) = \frac{P(A \cap R)}{P(R)} = \frac{\frac{1}{6}}{\frac{1}{2}} = \frac{1}{3}$$

43. Find the value of $P(R | S)$.

(a) $\frac{2}{3}$

(b) $\frac{3}{6}$

(c) $\frac{1}{3}$

(d) $\frac{4}{3}$

Correct Option - a: $\frac{2}{3}$

Explanation:

$$P(R \cap S) = \frac{2}{6}, P(R) = \frac{3}{6}$$

$$P(R|S) = \frac{P(R \cap S)}{P(R)} = \frac{\frac{2}{6}}{\frac{3}{6}} = \frac{2}{3}$$

44. The values of $P(A \cap R | S)$ and $P(A | S)$ are respectively

(a) $\frac{2}{3}$ and $\frac{1}{3}$

(b) $\frac{1}{3}$ and $\frac{2}{3}$

(c) $\frac{1}{6}$ and $\frac{3}{6}$

(d) $\frac{3}{6}$ and $\frac{1}{6}$

Correct Option - b: $\frac{1}{3}$ and $\frac{2}{3}$

Explanation:

$$P(A \cap R | S) = \frac{P(A \cap R \cap S)}{P(S)} = \frac{\frac{1}{6}}{\frac{3}{6}} = \frac{1}{3}$$

$$P(A|S) = \frac{P(A \cap S)}{P(S)} = \frac{\frac{2}{6}}{\frac{3}{6}} = \frac{2}{3}$$

45. Find the values of $P(A \cup R | S)$ and $P(R \cap S | A)$ respectively.

(a) $\frac{2}{5}$ and $\frac{1}{5}$

(b) $\frac{1}{5}$ and $\frac{2}{5}$

(c) 1 and $\frac{1}{2}$

(d) $\frac{1}{2}$ and 1

Correct Option - c: 1 and $\frac{1}{2}$

Explanation:

$$P(A \cup R | S) = \frac{P((A \cup R) \cap S)}{P(S)} = \frac{\frac{3}{6}}{\frac{3}{6}} = 1$$

$$P(R \cap S | A) = \frac{P(R \cap S \cap A)}{P(A)} = \frac{\frac{1}{6}}{\frac{2}{6}} = \frac{1}{2}$$

46. The revenue, R as a function of x can be expressed as

(a) $15x - \frac{x^2}{3000}$

(b) $15 - \frac{x^2}{3000}$

(c) $15x - \frac{1}{3000}$

(d) $15x - \frac{x}{3000}$

Correct option – a: $15x - \frac{x^2}{3000}$

Explanation:

The revenue function R(x) is given by

$$R(x) = p(x) \times x = \left(15 - \frac{x}{3000}\right)x$$

$$\Rightarrow R(x) = 15x - \frac{x^2}{3000}$$

47. The range of x is

(a) [0, 24000]

(b) [24000, 36000]

(c) [0, 36000]

(d) [12000, 24000]

Correct option – c: [0, 36000]

Explanation:

As the number of participants can be up to 36000.

So, the range of x is [0, 36000].

48. The value of x for which the revenue is maximum, is

(a) 20000

(b) 22500

(c) 21000

(d) 25000

Correct option – b: 22500

Explanation:

$$\text{Since, } R(x) = 15x - \frac{x^2}{3000}$$

$$\Rightarrow R'(x) = 15 - \frac{x}{1500}$$

For maxima/minima, $R'(x) = 0$

$$\Rightarrow 15 - \frac{x}{1500} = 0$$

$$\Rightarrow x = 22500$$

Again differentiating, we get

$$R'(x) = -\frac{1}{1500} < 0$$

At $x = 22500$, $f''(x) < 0$.

Hence, $x = 22500$ is the point of maxima.

49. If the revenue is maximum, the price of the ticket is

- (a) Rs. 5.5
- (b) Rs. 6
- (c) Rs. 7.5
- (d) Rs. 8

Correct option – c: Rs. 7.5

Explanation:

The revenue will be maximum at $x = 22500$

Therefore, price of a ticket is

$$15 - \frac{22500}{3000} = \text{Rs. } 7.5$$

50. How many students must participate so that the revenue is maximized?

- (a) 21000
- (b) 21500
- (c) 22000
- (d) 22500

Correct option – d: 22500

Explanation:

Number of students will be equal to the number of tickets sold.

Therefore, required number of students = 22500.